

Fusarium Wilt of Canary Island Date Palm¹

Monica L. Elliott²

Summary

- As the name implies, Fusarium wilt of Canary Island date palm is primarily observed on *Phoenix canariensis* (Canary Island date palm).
- The disease is caused by the fungus *Fusarium oxysporum* f. sp. *canariensis*.
- The leaf symptoms include a one-sided death, wherein the leaflets on only one side of the rachis are desiccated or dead. This is often accompanied by a reddish-brown or dark-brown streak on the petiole and rachis on the same side as the desiccated or dead leaflets. Eventually, the entire leaf dies.
- The disease symptoms normally appear first on the oldest (lowest) living leaves, and then progressively move upward in the canopy until the palm is killed.
- The only other disease that these symptoms could be confused with is petiole/rachis blight.
- Transmission of the fungus from palm to palm is primarily through contaminated pruning tools.
- There is no cure for this lethal disease. Fungicides have not been effective against Fusarium wilt.
- Since there is no cure, disease management is aimed at disease prevention. A new or disinfected pruning tool should be used for pruning leaves from each individual Canary Island date palm.
- A laboratory diagnosis using molecular techniques is required to confirm the Fusarium wilt pathogen.

Introduction

Unlike most fungal diseases of palms, this disease is very host specific, with the primary host being *Phoenix canariensis* (Canary Island date palm). It is caused by *Fusarium oxysporum* f. sp. *canariensis*. The name “Fusarium wilt of Canary Island date palm” was given to this disease to distinguish it from three other Fusarium wilt diseases that occur on palms. Fusarium wilt of Canary Island date palm was first documented in the United States in the 1970s in California, but it was not documented in Florida until 1994. It now occurs throughout the state.

The fungus causes a vascular wilt. Specifically, it obstructs the xylem (water-conducting) tissue, which results in desiccation and death, as described below.

The fungus produces short-lived spores (macroconidia and microconidia). It also produces chlamydospores, spores that live in the soil and plant tissue for long time periods (usually years). Because most of the root system is left behind after a palm dies from this disease, this mass of roots may act as a reservoir for the fungus for a long time.

Experimental and observational evidence suggests that other *Phoenix* species may be susceptible to *Fusarium oxysporum* f. sp. *canariensis*, including *Phoenix dactylifera* (date palm), *Phoenix reclinata* (Senegal date palm), and *Phoenix sylvestris* (wild date palm), but actual cases in the landscape or field nursery in Florida are extremely rare when compared to cases associated with *Phoenix canariensis*.

1. This document is PP-215, one of a series of the Plant Pathology Department, UF/IFAS Extension. Original publication date January 2006. Major revision August 2012; minor revision June 2013; Major revision October 2015 and June 2016. Reviewed March 2019. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.

2. Monica L. Elliott, professor, Plant Pathology Department, Fort Lauderdale Research and Education Center; UF/IFAS Extension, Gainesville, FL 32611.

The second *Fusarium* wilt disease of palms that occurs in Florida is “*Fusarium* wilt of queen and Mexican fan palm,” which primarily occurs on *Syagrus romanzoffiana* (queen palm) and *Washingtonia robusta* (Mexican fan palm). This disease is caused by *Fusarium oxysporum* f. sp. *palmarum*. See <http://edis.ifas.ufl.edu/pp278> for more information on this disease.

Palm Anatomy

To understand the symptoms described below, the terminology regarding palm anatomy must be defined. Phoenix species are an example of a pinnate or feather-leaf palm. The entire leaf is often called a frond. The leaf consists of a blade, a petiole, and a leaf base. The leaf base is the broad section that attaches to the trunk. The petiole is the “stem” of the leaf from the leaf base to the point where leaflets begin to attach to it. On a feather-leaf palm, the extension of the petiole into the leaf blade where all the leaflets attach is called the rachis. The leaflets are often referred to as pinnae.

Symptoms

Although the name “*Fusarium* wilt” implies that the primary symptom is a wilt, vascular wilts in palms look different from wilt diseases of hardwood trees. Instead of limp leaves or leaflets, they cause leaf desiccation and eventually leaf death symptoms. As with any disease, a progression of symptoms is observed with *Fusarium* wilt. Initial symptoms normally occur on the lowest (oldest) living leaves (Figure 1). Leaflets, either all of them or only a portion, and on only one side of the rachis, are discolored, usually a shade of brown because of desiccation or death. The leaflets on the opposite side of the rachis are healthy. This very distinctive symptom is often referred to as a “one-sided wilt” or “one-sided death” (Figure 2). Eventually, the leaflets on the other side of the rachis turn brown also, and the entire leaf dies.

A reddish-brown or dark-brown streak or stripe can be observed on the petiole and rachis of the affected frond, initially on the same side where the first dead leaflets appeared (Figure 3), and usually beginning at or near the leaf base (Figure 2). This streak may run the full length of the petiole and rachis or just a portion of it. Internal discoloration can be observed in cross section (Figure 4). This internal discoloration is caused by the fungus infesting the xylem tissue.

The disease symptoms normally begin on the lowest (oldest) leaves and then move up the canopy, progressively killing younger and younger leaves. The palm may die

relatively quickly (within months), but it usually lingers for as long as a year. The reason for this disparity is not known.



Figure 1. Overview of canopy of Canary Island date palm with *Fusarium* wilt. The black arrows point to leaves that are dying. Credits: M. L. Elliott



Figure 2. One-sided death of leaflets is typical of *Fusarium* wilt. Credits: T.K. Broschat and M. L. Elliott

Possible symptom variations include the disease initially starting mid-canopy rather than at the bottom of the canopy, perhaps because flower or fruit stalks were trimmed with an infested pruning tool. For some leaves, the leaflets may begin turning brown at the leaf tip and on both sides at once, rather than just on one side. It has also been noted that palm weevils in the bud of *Fusarium* wilt-stressed palms hasten the palm’s decline and may complicate determining the cause.



Figure 3. The reddish-brown stripe on the leaf petiole is on same side that exhibited one-sided death of leaflets from Fusarium wilt. Credits: N. A. Harrison and M. L. Elliott



Figure 4. Cross section of petiole shown in Figure 3. Note the internal discoloration is just on one side. Credits: M. L. Elliott

Diagnosis

Field diagnosis can often be made based on the symptoms described above, especially if the symptoms are typical ones. But symptoms of Fusarium wilt look the same as petiole/rachis blight. The latter is not a true vascular wilt and usually does not kill the palm, but the resulting individual leaf symptoms are very similar on *Phoenix* species. See the fact sheet about this disease at <http://edis.ifas.ufl.edu/pp145>. It is also possible for a Canary Island date palm to have both diseases. Therefore, a laboratory diagnosis is required to confirm Fusarium wilt.

To confirm the field diagnosis, a laboratory must isolate a *Fusarium oxysporum*-like fungus from the affected leaf

tissue and then conduct a molecular test on the isolated culture to confirm it is *f. sp. canariensis*. Contact your local county Extension office or the Florida Extension Plant Diagnostic Clinic (<http://www.plantclinic.org>) for complete details about correct sample submission procedures and cost of a laboratory diagnosis. The best tissue to send for a diagnosis is the petiole or rachis from a leaf exhibiting the typical symptoms of one-sided wilt. A dead leaf is not acceptable.

Molecular confirmation of the isolated fungus is critical. There are numerous *Fusarium oxysporum* isolates that do not cause disease but may be associated with palm material. Unfortunately, it is not possible to separate pathogenic isolates from nonpathogenic isolates based on their appearance in culture. Thus, isolating a fungus that looks like *F. oxysporum* does not confirm the field diagnosis. The fungus growing on the culture plate must be subjected to a molecular test that is specific for *F. oxysporum f. sp. canariensis*.

If the field symptoms appear typical for Fusarium wilt, but Fusarium is not isolated, then the symptoms may be an expression of petiole/rachis blight. Many of the petiole/rachis blight pathogens are not culturable from plant tissue. The next step would be to obtain new samples from the diseased palm or similarly diseased palm and resubmit to the clinic to confirm presence or absence of petiole/rachis blight pathogens. Be sure to discuss the problem with the clinic staff before resubmitting samples. If no symptomatic leaves are available, wait until new symptoms develop before resubmitting.

Disease Management

There are no cures for this disease. Fungicides have been evaluated, but none have been effective. Disease management is based solely on disease prevention.

The fungus is most likely moved between distant locations by plant material, both live plants and probably seed. This is probably how the disease was first introduced (and reintroduced) into Florida. Unpublished research from both Florida and Australia has demonstrated the fungus can colonize palm seeds. This makes it critical for palm nurseries to know that their seed and stock plants are coming from a disease-free source. Likewise, nursery growers, landscape architects, and installation companies should examine and select Canary Island date palms while they are still in the field or on site at a container nursery *and* make this inspection *prior* to any trimming of older leaves so the plants can be inspected for symptoms. Then, they should either monitor the trimming of these palms to ensure they

are trimmed correctly or trim them at the installation site using the correct procedure discussed below.

Once the disease is established in a landscape or nursery, the disease appears to primarily be transmitted from palm to palm via leaf pruning equipment (Figures 5 and 6). The fungal pathogen is in the vascular tissue of the leaf. Equipment used to remove leaves from an infected Canary Island date palm (chain saw, lopper, pruning shear, handsaw, etc.) will have fungal-infested leaf material (wood dust, plant sap) on the blades. If that equipment is not cleaned and disinfected, the next Canary Island date palm (and perhaps other *Phoenix* species) pruned by this equipment will be exposed to this fungal-infested leaf material.



Figure 5. The two Canary Island date palms in the foreground are dying from Fusarium wilt. The pathogen was likely moved from one to the other by pruning tools.

It is important to note that a palm could be infected but appear healthy (symptomless) because the disease has not developed to the point that leaf symptoms are being expressed. Unfortunately, it is not known how long this infected but symptomless stage of the disease lasts. It could be 6–18 months. Thus, the fungus could still be moved from palm to palm by pruning infected but symptomless leaves!

This means Canary Island date palms should be pruned with either new pruning tools or clean, disinfected tools, both in the landscape and nursery. This applies to all Canary Island date palms, diseased and healthy. **Either a new pruning tool or a clean, disinfected pruning tool should be used for each individual palm.** Pruning should be limited to once each year. Furthermore, pruning should be restricted to removal of *only* dead or dying leaves. Severe pruning, such as “hurricane cuts” or “pineapple cuts,” weakens trees and increases the risk of pathogen transmission. Pruning should be viewed as a risk factor for

Fusarium wilt disease transmission and not as a benefit to the Canary Island date palm.

Table 1 is a list of materials that can be used as disinfecting agents. Brush the tool blades clean of debris before placing in the disinfectant solution. For chain saws, it is recommended that they be taken apart and both the chain and bar soaked. By having multiple pruning tools, one tool can be soaking in the disinfectant solution in a bucket while the other tool is used for pruning. Alternatively, a newly developed pruning saw disinfecting tube with disinfectant solution can be attached to your belt (www.bladebath.com). The disinfectant solution should be replaced at least every 10 trees or every 2 hours. Rinse tools with clean water before pruning.

The following is a management suggestion: For the initial installation of a Canary Island date palm, it is normal to remove a large number of fronds. Using a handsaw would be time consuming, so either use a new chainsaw or disinfect the chainsaw between each palm. After installation, the number of leaves pruned should be limited as only dead leaves should be removed. This can be accomplished with a new pruning tool for each palm.

While this is an extraordinary measure, it is inexpensive disease prevention management for extremely valuable palms. A mature Canary Island date palm that has died from Fusarium wilt is expensive to remove and expensive to replace. It is certainly more economical to prevent the disease than deal with the deadly consequences, especially if there are multiple Canary Island date palms in the landscape (Figure 6). Using a new pruning tool or disinfecting the pruning tool is inexpensive in comparison to a devastated landscape.



Figure 6. Multiple *Phoenix canariensis* dying from Fusarium wilt. The disease is moved from palm to palm by infested pruning tools. Credits: H. M. Donselman

Again, this is a disease without a cure. In a nursery situation, diagnosis of this disease effectively destroys the crop as the plants are unsaleable and must be destroyed. In the landscape situation, the palm eventually dies and must be removed. In both situations, but especially the nursery situation, the diseased palm(s) should be removed and destroyed immediately. Be very careful not to distribute potentially fungal-infested soil or potting mix in the nursery. Surrounding Canary Island date palms should be monitored closely because it is quite likely they could be infected.

The diseased palm should be incinerated or placed in a landfill. If this is not possible (as is the case in many Florida counties), then be sure the palm is not chipped and recycled for mulch in the landscape. Instead, chipped material should be placed in a compost pile. Chain saws and other tools used for removal must be brushed free of plant material and disinfected as described previously.

Chlamydospores of *Fusarium oxysporum* f. sp. *canariensis* can survive for long periods in the soil and certainly will survive in any infected root mass left behind when a diseased palm is removed. Experimental and observational evidence has shown the potential for infecting palms, especially seedling palms, with this fungus via the root system. Therefore, it would not be prudent to plant a Canary Island date palm back into a site where *Fusarium* wilt killed the previous palm(s). It is not known how far away from the potentially fungal-infested site one must move before it is safe to plant a Canary Island date palm.

Selected References

Feather, T. V. 1982. "Occurrence, Etiology and Control of Wilt and Dieback of *Phoenix canariensis* in California." Ph.D. diss. University of California, Riverside, CA.

Feather, T. V., D. E. Munnecke, and J. B. Carpenter. 1989. "The Occurrence of *Fusarium oxysporum* on *Phoenix canariensis*: A Potential Danger to Date Production in California." *Plant Disease* 73 (1): 78–80.

Plyler, T. R., G. W. Simone, D. Fernandez, and H. C. Kistler. 1999. "Rapid Detection of the *Fusarium oxysporum* Lineage Containing the Canary Island Date Palm Wilt Pathogen." *Phytopathology* 89 (5): 407–413.

Simone, G. W. 1998. *Prevention and Management of Palm Diseases in Florida's Landscapes*. Plant Pathology Mimeo 98-4. Gainesville: University of Florida Cooperative Extension Service.

Simone, G. W. 2004. "Fusarium Wilt." In *Compendium of Ornamental Palm Diseases and Disorders*, edited by M. L. Elliott, T. K. Broschat, J. Y. Uchida, and G. W. Simone, 17–25. St. Paul, MN: American Phytopathological Society.

Simone, G. W., and G. Cashion. 1996. *Fusarium Wilt of Canary Island Date Palms in Florida*. Plant Pathology Fact Sheet PP-44. Gainesville: University of Florida Cooperative Extension Service.

Smith, D. I., I. W. Smith, and P. Clements. 2003. *Fusarium Wilt of Canary Island Date Palm*. Forests Fact Sheet, Victoria, Australia: Department of Sustainability and Environment.

Summerell, B. A., H. C. Kistler, and L. V. Gunn. 2001. "Fusarium Wilt of *Phoenix canariensis* Caused by *Fusarium oxysporum* f. sp. *canariensis*." In *Fusarium: Paul E. Nelson Memorial Symposium*, edited by B. A. Summerell, J. F. Leslie, D. Backhouse, W. L. Bryden, and L. W. Burgess, 263–270. St. Paul, MN: American Phytopathological Society.

Table 1. Suggested materials and soaking times for disinfecting pruning tools

Material*	% solution	Soaking time
Household bleach (e.g., Clorox®)	25% (1 part bleach + 3 parts water)	5 minutes minimum
Pine oil cleaner (e.g., Pine Sol®)	25% (1 part cleaner + 3 parts water)	5 minutes minimum
Rubbing alcohol (70% isopropyl)	50% (1 part alcohol + 1 part water)	5 minutes minimum
Denatured ethanol (95%)	50% (1 part alcohol + 1 part water)	5 minutes minimum

*The above materials were shown to be effective in eliminating the fungus from the wood dust or palm sap trapped on pruning tools (Simone 1998). It is suggested that the solution be replaced after 10 trees or every 2 hours. Rinse the tool with fresh water after soaking. Other potential disinfectants include trisodium phosphate or quaternary ammonium salts. The latter is recommended at a 5% solution, soaking for 5 minutes (Smith, Smith, and Clements 2003).