

CRISPR Gene Editing in Strawberry¹

Seonghee Lee, Cheolmin Yoo, Yoon Jeong Jang, Kevin Folta, and Vance M. Whitake²

Recent technology has been developed to precisely engineer genes for traits of interest. This approach is known as “CRISPR” gene editing. Gene editing is distinct from other forms of genetic engineering, such as transgenic technology, often colloquially referred to as “GMO”. With gene editing, the final product can match that obtained by conventional plant breeding, but in a much shorter timeframe. CRISPR has been applied in many agronomic crops and is poised to make contributions in strawberry. We anticipate that, over the next decade, CRISPR and other gene editing techniques will be used to rapidly develop elite strawberry varieties with improved disease resistance, fruit quality, and other valuable attributes.

What is CRISPR gene editing?

One of the great disappointments in the pursuit of improved varieties is the discovery of a new advanced selection that would be valued by the industry except for one critical flaw. Gene editing technology can be used to almost surgically adjust the gene or genes behind that deleterious trait. CRISPR (Clustered Regularly Interspaced Short Palindromic Repeats) is one form of gene editing that can be used to precisely modify a gene of interest without otherwise compromising the favorable traits of an elite variety (Rani et al. 2016; Bortesi and Fischer 2015). For example, instead of breeding for many years to move a disease resistance gene from a wild strawberry into a modern strawberry, gene editing allows a direct introduction of the genetic information. Think of it as a “cut and

paste” mechanism. This is particularly useful for cultivated strawberries because they are genetically complex, making conventional breeding difficult. The UF/IFAS strawberry breeding program has identified several important gene regions controlling disease resistance traits that are directly relevant to Florida growers. By using CRISPR technology, these genes or gene variants can be moved into desirable genetic backgrounds that can be further moved via conventional crossing in later generations. We can utilize established DNA marker-assisted breeding tools to track the edited genes in subsequent generations, adding to the speed of new variety development. Based on policy discussions, it is possible that the first-generation plants containing the edited genes will not require extensive regulation, and these tools will be extremely valuable in the long-term efforts of every strawberry breeding program.

Does CRISPR gene editing = transgenic technology?

“Transgenic technology” refers to the transfer of a genetic material from one species to another. CRISPR, on the other hand, can be used to precisely change DNA sequence, switching it from one naturally occurring variant to another naturally occurring variant. Using this new technique, we can cut a strawberry’s genome at a desired location so that existing genes can be removed or added. Early indications suggest that gene editing should be regulated like conventionally bred crops, as the final product can simply match what may be done by conventional breeding. Recently,

1. This document is HS1315, one of a series of the Horticultural Sciences Department, UF/IFAS Extension. Original publication date February 2018. Revised December 2021. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Seonghee Lee, assistant professor; Cheolmin Yoo, biological scientist; Yoon Jeong Jang, Postdoctoral Associate; Kevin Folta, professor; and Vance M. Whitaker, associate professor; Horticultural Sciences Department, UF/IFAS Gulf Coast Research and Education Center; UF/IFAS Extension, Gainesville, FL 32611.

the USDA announced that it would not regulate a new mushroom developed using CRISPR (Waltz 2016; Hoffman 2021). Countries like Sweden and Argentina have made similar proclamations, indicating that the finished CRISPR-edited varieties do not fall under certain regulations because they do not contain “foreign” DNA. We published another article about CRISPR gene editing in crop improvement (<https://edis.ifas.ufl.edu/publication/HS1334>).

Development of a CRISPR System in Strawberry

While the finished varieties do not contain DNA from other organisms, the process introduces genetic information that orchestrates the desired genetic change. The first step is to develop and optimize a tissue culture and transformation system (protocols to introduce foreign DNA to new plants) for UF strawberry lines, so that new plants can be regenerated from cells containing introduced DNA. However, just as each cultivar has different traits and qualities, they also behave differently with respect to introduction of new genes.

The major strawberry cultivars grown in Florida such as ‘Florida Radiance’, Sweet Sensation® ‘Florida127’ and ‘Florida Brilliance’ were used for tissue culture optimization. As shown in Figure 1, callus induction was tested with different strawberry tissues, and embryogenic callus growth was most vigorous on stolons (runners) and petioles. To identify the optimal conditions for shoot and root regeneration for ‘Florida Radiance’, Sweet Sensation® ‘Florida127’ and ‘Florida Brilliance’, explants were grown on a range of media with varying compositions of plant growth regulators.

C:\LyraEDISServlet?command=getScreenImage&id=15623809

About one inch of petiole or stolon from the leaf-end or shoot-end, respectively, were collected from greenhouse grown plants and used for the tissue culture process. Optimal conditions for tissue culture medium, nutrient, and hormone were tested for the UF accessions. The runner (stolon) produced calli more vigorously than petiole segments. It takes about 14 weeks for ‘Florida Radiance’, Sweet Sensation® ‘Florida127’ and ‘Florida Brilliance’ to develop from embryogenic callus to young plantlet in rooting media (Figure 2). ‘Florida Brilliance’ produced more regenerated plants than Sweet Sensation® ‘Florida127’.



Figure 1. Test for callus induction with different tissues of Sweet Sensation® ‘Florida 127’. Leaf tissues, meristem tissues from crowns, stolons, and petioles were harvested from greenhouse-grown plants that were six to eight months old.

Credits: Cheolmin Yoo




Figure 2. Progression of shoot and root regeneration for Sweet Sensation® ‘Florida 127’ and ‘Florida Brilliance’ (FL 13.26-134). First image from left: shoots and embryos are generated from the callus of the explants; second image: expanded shoots; third image: elongated shoot clusters; fourth image: plantlets in rooting media.

Credits: Cheolmin Yoo

No “Foreign” DNA Sequences

Once the genetically engineered gene product is ready for CRISPR gene editing, transformation, where the new genetic material is delivered to a single strawberry cell, is the first step in the genetic engineering process. For DNA delivery, *Agrobacterium tumefaciens*-mediated transformation is widely used for CRISPR gene editing. *Agrobacterium tumefaciens* is a widespread, naturally occurring soil bacterium that causes crown gall in many plant species and has the ability to introduce new genetic material into plant cells (Gelvin 2003). This bacterium works as a natural genetic engineer and is used in labs for plant transformation. Gene-edited plants using *Agrobacterium*-mediated transformation will contain foreign bacterial DNA sequences. It is not an easy process to remove the bacterium-derived DNA sequences through breeding.

In contrast, we are currently using a biolistic particle bombardment method in which DNA-coated metal particles are delivered to the plant cells using a “gene gun” (Figure 3).




Figure 3. (A) General procedure of transformation using the metal particle bombardment system. (B) Transient expression of synthetic green fluorescent protein (GFP) in ‘Sweet Sensation®’ strawberry. (C) CRISPR precisely knocked out the chlorophyll gene, *Phytoene Desaturase (PDS)*, in Sweet Sensation® ‘Florida 127’. GFP signal in strawberry callus showing that transformation was successfully done (a-b), CRISPR mutant shoot lines with albino phenotype (c-f)
Credits: UF/IFAS and Jin-Hee Kim

This method can be applied to a wide range of cell and tissue types, and there is no need for bacterial DNA insertion. Later, the gene edited tissues can be regenerated to mature plants using the tissue culture protocol outlined above. The GFP was delivered to cells using a hand-held “gene gun”. When GFP is successfully delivered into cells, green fluorescence can be observed under a UV-microscopy (Figure 3B). Since we already understand the sequence of a gene involved in chlorophyll synthesis called the “*Phytoene Desaturase (PDS)*” gene, we have used CRISPR to interfere with the function of this gene as a way to test whether our strawberry CRISPR system works. Figure 2 shows that CRISPR precisely knocked out the chlorophyll gene in Sweet Sensation® (Figure 3C).

Recently, the development of gene editing using protoplasts and regeneration of plants from protoplasts has been demonstrated in other plant species (Woo et al. 2015; Kan-chiswamy 2016; Dutt et al. 2015). This method is known as a completely “DNA-free” gene editing system. Briefly, the protein/genetic material complex for gene editing will be assembled in vitro and the complex mixed with strawberry protoplast isolated from embryogenic calli and polyethylene glycol, which allows direct transfer by endocytosis into protoplasts. The gene edited with protoplasts is cultured (cell suspension culture) into calli, and mature plants can be regenerated using the tissue culture protocol outlined above. We are currently developing the protocol of CRISPR gene editing with strawberry protoplast (Figure 4).




Figure 4. Isolation of strawberry protoplasts from in-vitro cultured Sweet Sensation® ‘Florida 127’. Protoplast released 15 hours after enzyme digestion (A), and tissue debris was filtered for purifying protoplasts (B). The freshly isolated strawberry protoplast were transformed with CRISPR constructs (C).
Credits: undefined

Gene Editing for Disease Resistance

The UF strawberry breeding program provides a direct commercialization path for disease-resistant varieties to reach Florida growers. New varieties with better resistance will bring increased profitability to the Florida strawberry industry. The Florida Strawberry Growers Association estimates that diseases cost the Florida industry at least \$15 million each year. In the last five years, the UF strawberry breeding program has identified regions of strawberry chromosomes that control fruit flavor and multiple disease resistance (Barbey et al. 2021; Chandra et al. 2021; Nelson et al. 2021; Noh et al. 2018; Oh et al. 2021; Oh et al. 2020; Oh et al. 2019; Salinas et al. 2020). Efforts are ongoing to narrow these chromosomal regions down and identify the exact gene sequences that provide these disease resistances, with a recent emerging strawberry disease, *Neopestalotiopsis*, resistance as the first priority. Our goal is to add *Neopestalotiopsis* resistance to ‘Florida Brilliance’, Sweet Sensation® ‘Florida127’, and new cultivars such as ‘Florida Medallion’, which are currently highly susceptible to this disease. Evaluations of the gene-edited lines will be performed in concert with crosses to integrate the changes into other major varieties and advanced selections with conventional hybridization.

References

- Barbey CR et al. 2021. “Genetic analysis of methyl anthranilate, mesifurane, linalool, and other flavor compounds in cultivated strawberry (*Fragaria × ananassa*)” *Frontiers in Plant Science*. 12: 615749.
- Bortesi, L. and R. Fischer. 2015. “The CRISPR/Cas9 system for plant genome editing and beyond.” *Biotechnol Adv*. 33(1): 41-52.
- Chandra S et al. 2021. “Comparative transcriptome analysis to identify candidate genes for *FaRCg1* conferring resistance against *Colletotrichum gloeosporioides* in cultivated strawberry (*Fragaria × ananassa*)” *Frontiers in Genetics*. 12: 730444.
- Dutt, M. et al. 2015. “Transgenic Citrus Expressing an Arabidopsis NPR1 Gene Exhibit Enhanced Resistance against Huanglongbing (HLB; Citrus Greening)” *PLoS One* 10(9):e0137134.
- Gelvin, S.B. 2003. “Agrobacterium-mediated plant transformation: the biology behind the “gene-jockeying” tool.” *Microbiol Mol Biol Rev*. 67(1):16-37.
- Hoffman, N. E. “Revisions to USDA biotechnology regulations: The SECURE rule.” *Proceedings of the National Academy of Sciences of the United States of America*.
- Kanchiswamy, C.N. 2016. “DNA-free genome editing methods for targeted crop improvement.” *Plant Cell Rep*. 35(7): 1469-1474.
- Mangandi, J. et al. 2017. “Pedigree-Based Analysis in a Multiparental Population of Octoploid Strawberry Reveals QTL Alleles Conferring Resistance to *Phytophthora cactorum*.” *G3* 7(6): 1707-1719.
- Nelson JR et al. 2021. “Discovery of three loci increasing resistance to charcoal rot caused by *Macrophomina phaseolina* in octoploid strawberry.” *G3* 11:jkab037.
- Noh Y-H et al. 2018. “High-throughput marker assays for *FaRPC2*-mediated resistance to *Phytophthora crown rot* in octoploid strawberry.” *Molecular Breeding* 38:104.
- Oh Y et al. 2021. “Genomic characterization of the fruity aroma gene, *FaFAD1*, reveals a gene dosage effect on γ -decalactone production in strawberry (*Fragaria × ananassa*)” *Frontiers in Plant Science* 12: 639345.
- Oh Y, et al. 2020. “Development of subgenome-specific markers for *FaRXf1* conferring resistance to bacterial angular leaf spot in allo-octoploid strawberry.” *Int J Fruit Sci*. 1-13.
- Oh Y, et al. 2019. “The strawberry DNA testing handbook.” *HortScience* 54: 2267-2270.
- Rani, R., et al. 2016. “CRISPR/Cas9: a promising way to exploit genetic variation in plants.” *Biotechnol Lett*.
- Roach, J.A., et al. 2016. “*FaRXf1*: a locus conferring resistance to angular leaf spot caused by *Xanthomonas fragariae* in octoploid strawberry.” *Theor Appl Genet*. 129(6): 1191-1201.
- Salinas N, Fan Z, Peres N, Lee S, Whitaker VM (2020) *FaRCa1* confers moderate resistance to the root necrosis form of strawberry anthracnose caused by *Colletotrichum acutatum* *HortScience* 55:693-698
- Waltz, E. 2016. “Gene-edited CRISPR mushroom escapes US regulation.” *Nature* 532(7599): 293.
- Woo, J.W., et al. 2015. “DNA-free genome editing in plants with preassembled CRISPR-Cas9 ribonucleoproteins.” *Nat Biotechnol*. 33(11): 1162-1164.